

July 1997
UM-P-97/48
RCHEP-97/08

An alternative $SU(4) \otimes SU(2)_L \otimes SU(2)_R$ model

R. Foot

*School of Physics
Research Centre for High Energy Physics
The University of Melbourne
Parkville 3052 Australia*

Abstract

A simple alternative to the usual Pati-Salam model is proposed. The model allows quarks and leptons to be unified with gauge group $SU(4) \otimes SU(2)_L \otimes SU(2)_R$ at a remarkably low scale of about 1 TeV. Neutrino masses in the model arise radiatively and are naturally light.

In the standard $SU(2)_L \otimes U(1)_Y$ model of electroweak interactions[1] quarks and leptons have many similarities. For example, there are three generations of quarks and three generations of leptons, with 2 quarks and 2 leptons in each generation. Furthermore both left-handed quarks and left-handed leptons transform as $SU(2)_L$ doublets while both right-handed quarks and right-handed leptons are $SU(2)_L$ singlets.

The similarity of quarks and leptons may be due to a spontaneously broken exact symmetry. This symmetry if it exists could be either continuous[2] or discrete[3]. In the case where it is discrete, unification of quarks and leptons can occur at very low scales of around a TeV[3, 4]. If quarks and leptons are related by a continuous Pati-Salam type gauge symmetry it is usually assumed that this symmetry is broken at a very high scale $M \geq 10^{11} \text{ GeV}$ which means that the idea cannot be tested directly in any conceivable experiment. However if there is no left-right symmetry then it is nevertheless still possible that the standard model is a remnant of a gauge model with Pati-Salam gauge symmetry broken at a relatively low scale of 1000 TeV or even less[5, 6, 7]. The purpose of this letter is to point out that there exists an interesting alternative Pati-Salam type model which can be broken at a much lower scale of the order of a TeV.

The gauge symmetry of the model is

$$SU(4) \otimes SU(2)_L \otimes SU(2)_R. \quad (1)$$

Under this gauge symmetry the fermions of each generation transform in the anomaly free representations:

$$Q_L \sim (4, 2, 1), \quad Q_R \sim (4, 1, 2), \quad f_L \sim (1, 2, 2). \quad (2)$$

The minimal choice of scalar multiplets which can both break the gauge symmetry correctly and give all of the charged fermions mass is

$$\chi_L \sim (4, 2, 1), \quad \chi_R \sim (4, 1, 2), \quad \phi \sim (1, 2, 2). \quad (3)$$

These scalars couple to the fermions as follows:

$$\mathcal{L} = \lambda_1 \bar{Q}_L (f_L)^c \tau_2 \chi_R + \lambda_2 \bar{Q}_R f_L \tau_2 \chi_L + \lambda_3 \bar{Q}_L \phi \tau_2 Q_R + \lambda_4 \bar{Q}_L \phi^c \tau_2 Q_R + H.c., \quad (4)$$

where the generation index has been suppressed and $\phi^c = \tau_2 \phi^* \tau_2$. Under the $SU(3)_c \otimes U(1)_T$ subgroup of $SU(4)$, the 4 representation has the branching rule, $4 = 3(1/3) + 1(-1)$. We will assume that the $T = -1, I_{3R} = -1/2$ ($I_{3L} = 1/2$) components of χ_R (χ_L) gain non-zero Vacuum Expectation Values (VEVs) as well as the $I_{3L} = I_{3R} = -1/2$ and $I_{3L} = I_{3R} = 1/2$ components of the ϕ . We denote these VEVs by $w_{R,L}, u_{1,2}$ respectively. In other words,

$$\begin{aligned} \langle \chi_R(T = -1, I_{3R} = -1/2) \rangle &= w_R, \quad \langle \chi_L(T = -1, I_{3L} = 1/2) \rangle = w_L, \\ \langle \phi(I_{3L} = I_{3R} = -1/2) \rangle &= u_1, \quad \langle \phi(I_{3L} = I_{3R} = 1/2) \rangle = u_2. \end{aligned} \quad (5)$$

We will assume that the VEVs satisfy $w_R > u_{1,2}, w_L$ so that the symmetry is broken as follows:

$$\begin{aligned}
& SU(4) \otimes SU(2)_L \otimes SU(2)_R \\
& \quad \downarrow \langle \chi_R \rangle \\
& SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \\
& \quad \downarrow \langle \phi \rangle, \langle \chi_L \rangle \\
& SU(3)_c \otimes U(1)_Q
\end{aligned} \tag{6}$$

where $Y = T - 2I_{3R}$ is the linear combination of T and I_{3R} which annihilates $\langle \chi_R \rangle$ (i.e. $Y\langle \chi_R \rangle = 0$). Observe that in the limit where $w_R \gg w_L, u_1, u_2$, the model reduces to the standard model. The VEV w_R breaks the gauge symmetry to the standard model subgroup. This VEV also gives large (electroweak invariant) masses to an $SU(2)_L$ doublet of exotic fermions, which have electric charges $-1, 0$. We will denote these exotic fermions with the notation E^-, E^0 . These exotic fermions must have masses greater than $M_Z/2$ otherwise they would contribute to the Z width. Observe that the right-handed chiral components of the usual charged leptons are contained in Q_R . They are the $T = -1, I_{3R} = 1/2$ components. The usual left-handed leptons are contained in f_L along with the right-handed components (CP conjugated) of E^0, E^- . It is instructive to write out the fermion multiplets explicitly. For the first generation,

$$Q_L = \begin{pmatrix} d_y & u_y \\ d_g & u_g \\ d_b & u_b \\ E^- & E^0 \end{pmatrix}_L, \quad Q_R = \begin{pmatrix} u_y & d_y \\ u_g & d_g \\ u_b & d_b \\ \nu & e \end{pmatrix}_R, \quad f_L = \begin{pmatrix} (E_R^-)^c & \nu_L \\ (E_R^0)^c & e_L \end{pmatrix}, \tag{7}$$

and similarly for the second and third generations. In the above matrices the first column of Q_L (f_L, Q_R) is the $I_{3L}(I_{3R}) = -1/2$ component while the second column is the $I_{3L}(I_{3R}) = 1/2$ component. The four rows of Q_L, Q_R are the four colours and the rows of f_L are the $I_{3L} = \pm 1/2$ components. Observe that the VEVs $w_L, u_{1,2}$ have the quantum numbers $I_{3L} = -1/2, Y = 1$ (or equivalently $I_{3L} = 1/2, Y = -1$). This means that the standard model subgroup, $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ is broken to $SU(3)_c \otimes U(1)_Q$ in the usual way (with $Q = I_{3L} + Y/2$).

Having established that the model is a phenomenologically viable extension to the standard model, we now comment on various features of the model.

Observe that the model has the rather unusual feature that the scalar multiplets required to break the gauge symmetry and give the fermions masses have precisely the same quantum numbers as the fermion multiplets of a generation [compare Eq.(2) and Eq.(3)].

In the model the ordinary neutrinos are naturally light. The neutrino masses vanish at tree-level given the particle content of the theory. The model does however have a light singlet neutrino, ν_R . This electroweak singlet occupies the $T = -1, I_{3R} = -1/2$ component of the Q_R multiplet. Note however that with the minimal Higgs content, this field cannot couple to the ordinary left-handed neutrinos (at tree level). Observe that Majorana neutrino masses arise from the $W_{L,R}$ gauge interactions at the one loop level. Assuming diagonal couplings and examining ν_e for definiteness, the Feynman diagram for the ν_e mass is shown in Figure 1. Calculating this finite 1-loop diagram we find that

$$m_\nu \simeq \frac{2g_R g_L}{(4\pi)^2} \left[\frac{g_R g_L u_1 u_2}{M_{W_R}^2} \right] \left[\frac{m_e m_d M_E}{M_E^2 - M_{W_L}^2} \right] \log \left(\frac{M_E^2}{M_{W_L}^2} \right), \quad (8)$$

where $g_{L,R}$ are the $SU(2)_{L,R}$ gauge coupling constants and we have assumed that $M_E^2 \ll M_{W_R}^2$. Clearly the neutrino masses are naturally light given that $M_{W_R}, M_E \gg m_e, m_d$.

The gauge interactions of the model conserve an unbroken baryon number symmetry. This baryon charge is defined as $B = B' + T$ where the B' charges of $Q_L, Q_R, \chi_{L,R}$ are 1 and the B' charges of f_L, ϕ are 0. The existence of the baryon number symmetry implies that protons and neutrons are absolutely stable in the model.

Because the right-handed charged leptons belong to the same multiplet as the right-handed quarks there will be gauge interactions of the form

$$\mathcal{L} = \frac{g_s}{\sqrt{2}} \bar{D}_R^i W'_\mu \gamma^\mu K'^{ij} l_R^j + H.c., \quad (9)$$

where $i, j = 1, \dots, 3$ are family indices, that is $D_R^1 = d_R, D_R^2 = s_R, D_R^3 = b_R, l_R^1 = e_R, l_R^2 = \mu_R, l_R^3 = \tau_R$ and W'_μ are electrically charged $2/3$ gauge bosons (which gain masses from χ_R at the first step of symmetry breaking). The matrix K' is a Cabibbo-Kobayashi-Maskawa type matrix. The most stringent bound on the symmetry breaking scale w_R is expected to arise from $K_L \rightarrow \mu^\pm e^\mp$ decays. The decay $\bar{K}^0 \rightarrow \mu^- e^+$ arises from a Feynman diagram with a T-channel exchange of a W' gauge boson. This diagram corresponds (after a Fierz rearrangement) to the effective four fermion Lagrangian density,

$$\mathcal{L}_{eff} = \frac{G_X}{\sqrt{2}} \bar{d} \gamma_\mu (1 + \gamma_5) s \bar{\mu} \gamma^\mu (1 + \gamma_5) e + H.c., \quad (10)$$

where $G_X = \sqrt{2} g_s^2 (M_{W'}) / 8 M_{W'}^2$. Using this effective Lagrangian it is straightforward to calculate the decay rate. We find,

$$\Gamma(\bar{K}^0 \rightarrow \mu^- e^+) \simeq \frac{G_X^2 f_K^2}{8\pi} M_K m_\mu^2, \quad (11)$$

where f_K is the K meson decay constant, M_K is the K meson mass and we have assumed that the mixing matrix K'_{ij} is approximately diagonal. Evaluating the above equation we find that

$$Br(\bar{K}^0 \rightarrow \mu^- e^+) \simeq 10^{-2} \left(\frac{TeV}{M_{W'}} \right)^4. \quad (12)$$

The current experimental bound, $Br(\bar{K}^0 \rightarrow \mu^\pm e^\mp) < 3.3 \times 10^{-11}$ [8], implies the limit

$$M_{W'} \gtrsim 140 \text{ TeV}, \quad (13)$$

assuming that the mixing matrix K'^{ij} is approximately diagonal. This bound is the most stringent bound on the model in the case where K'^{ij} is diagonal. However in the model there is no relationship between the charged lepton mass matrix and the quark masses. Indeed, in the model they are proportional to the VEVs of different scalar multiplets and have independent Yukawa couplings. One consequence of this is that the mixing matrix K'^{ij} connecting the right-handed quarks with the right-handed leptons is theoretically unconstrained (except of course, for the unitary requirement). For example, the W' could couple s_R predominately with τ_R (this possibility was discussed in the context of the usual Pati-Salam model in Ref.[6]). If this is the case then K_L decays do not give any stringent bounds on the model. In order to explore this scenario further, we will assume for definiteness that W' couples s_R with τ_R , d_R with e_R and b_R with μ_R . This corresponds to a K'^{ij} matrix of the form

$$K' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (14)$$

Of course it would not seem natural for the zero elements of this mixing matrix to be exactly zero. However we will assume that they are zero to illustrate a point. In the case of the usual Pati-Salam model with the ansatz Eq.(14), the most stringent bound on $M_{W'}$ comes from the W' contribution to the decay $\pi^+ \rightarrow e^+ \nu_L$ [6]. This process leads to a quite stringent bound of $M_{W'} \gtrsim 250 \text{ TeV}$ [6] for that model. This bound arises by calculating the interference term between the amplitudes arising from the standard model contribution and the Pati-Salam contribution. However the decay $\pi^+ \rightarrow e^+ \nu$ does not provide a stringent constraint for the alternative Pati-Salam model. There are two reasons for this. First, the W' mediates the decay $\pi^+ \rightarrow e^+ \nu_R$ (rather than $\pi^+ \rightarrow e^+ \nu_L$). Because the final state is distinct from the standard model process $\pi^+ \rightarrow e^+ \nu_L$, there will obviously be no interference term between the amplitudes of the two processes. Second, the W' of the alternative model only couples to right-handed

quarks and leptons. This means that the decay $\pi^+ \rightarrow e^+ \nu_R$ is helicity suppressed by a factor m_e^2/m_π^2 (which is also the case for the standard model contribution). In fact,

$$\frac{\Gamma(\pi^+ \rightarrow e^+ \nu_R)}{\Gamma(\pi^+ \rightarrow e^+ \nu_L)} \simeq \frac{G_X^2}{G_F^2}, \quad (15)$$

where G_F is the usual Fermi constant. The above contribution to $\pi^+ \rightarrow e^+ \nu$ decay leads to a violation of lepton universality and implies a small modification to the ratio R where $R \equiv \Gamma(\pi^+ \rightarrow e^+ \nu)/\Gamma(\pi^+ \rightarrow \mu^+ \nu)$. Using $\alpha_s(M_{W'}) \sim 1/10$, we find,

$$\frac{\delta R}{R} \simeq 4 \times 10^{-4} \left(\frac{TeV}{M_{W'}} \right)^4. \quad (16)$$

The theoretical prediction for R agrees within errors to the experimental measurement and thus Eq.(16) can be compared to the experimental error $\delta R/R \sim 0.003$ [8]. Clearly then, π^+ decay does not lead to any significant bound for the model.

A more stringent bound on the model [assuming the ansatz Eq.(14)] arises from the W' mediated rare B_d^0 decay, $\bar{B}_d^0 \rightarrow \mu^- e^+$. In the case of the usual Pati-Salam model with the ansatz Eq.(14), the bound $M_{W'} \gtrsim 16 TeV$ was derived in Ref.[6]. However in my model the bound arising from this process is much less stringent. The main difference is that the W' of the usual Pati-Salam model couples vectorially where as the W' of the alternative Pati-Salam model couples only to right-handed quarks and leptons and $\bar{B}_d^0$ decays will be helicity suppressed by a factor $\sim m_\mu^2/M_B^2$. In fact the width for the decay $\bar{B}_d^0 \rightarrow \mu^- e^+$ [assuming the ansatz Eq.(14)] is given by Eq.(11) with the replacement $K \rightarrow B$. Evaluating the resulting equation we find that

$$Br(\bar{B}_d^0 \rightarrow \mu^- e^+) \simeq 3 \times 10^{-6} \left(\frac{TeV}{M_{W'}} \right)^4. \quad (17)$$

The current experimental bound, $Br(\bar{B}_d^0 \rightarrow \mu^\pm e^\mp) < 6 \times 10^{-6}$ [8], implies the limit

$$M_{W'} \gtrsim 800 GeV. \quad (18)$$

If the W' gauge boson is light, then how large can the zero elements of K'_{ij} be? The most stringently constrained element is the $K'_{s\mu}$ entry [which is the K'_{22} element of Eq.(14)]. This entry is constrained to be $K'_{s\mu} \lesssim 10^{-4}$ if $M_{W'} \simeq 1 TeV$ given the experimental bound $Br(K_L \rightarrow \mu^\pm e^\mp) < 3.3 \times 10^{-11}$ [8].

Note that so-called vector lepto-quarks have been studied which have similar properties to the W' gauge bosons[9]. However it is usually assumed that vector lepto-quarks must couple to only one generation if they are to be light enough to be seen in collider experiments. One result of this paper is that it is possible to have light vector

lepto-quarks coupling chirally to all three quarks and leptons. Furthermore, the model provides a concrete renormalizable framework where vector lepto-quarks with chiral couplings are gauge fields and may thus be fundamental particles.

In addition to the exotic W' gauge bosons, the model contains $W_R^\pm$ and Z' gauge bosons. The exotic gauge boson mass matrix arises from the Lagrangian density terms (in the limit where $w_R \gg w_L, u_{1,2}$)

$$\mathcal{L} = (D_\mu \langle \chi_R \rangle)^\dagger D^\mu \langle \chi_R \rangle. \quad (19)$$

where the covariant derivative is given by

$$D_\mu = \partial_\mu + ig_s G_\mu^a \Lambda_a + ig_L W_{L\mu}^i \tau_L^i / 2 + ig_R W_{R\mu}^i \tau_R^i / 2, \quad (20)$$

where $a = 1, \dots, 15, i = 1, \dots, 3$ and $G_\mu^a, W_{L\mu}^i, W_{R\mu}^i$ ($\Lambda_a, \tau_L^i/2, \tau_R^i/2$) are the $SU(4), SU(2)_L, SU(2)_R$ gauge bosons (generators) respectively. By examining the exotic gauge boson mass matrix Eq.(19), it is possible to obtain the usual weak mixing angle, $\sin \theta_w \equiv e/g_L$, as a function of the couplings g_s, g_L, g_R . We find that

$$\sin^2 \theta_w(M_{W'}) = \frac{g_s^2(M_{W'}) g_R^2(M_{W'})}{g_s^2(M_{W'}) g_R^2(M_{W'}) + g_s^2(M_{W'}) g_L^2(M_{W'}) + \frac{2}{3} g_L^2(M_{W'}) g_R^2(M_{W'})}. \quad (21)$$

Assuming that $\sin^2 \theta_w(M_{W'}) \simeq 1/4$ which is appropriate for $M_{W'} \sim 1 \text{ TeV}$, Eq.(21) implies that $g_R(M_{W'}) \simeq g_L(M_{W'})/\sqrt{3}$. Also it is easy to show that $M_{W'} \simeq \sqrt{2/3} M_{Z'} \simeq (g_s/g_R) M_{W_R}$. Furthermore the Z' gauge boson couples to fermions via the interaction Lagrangian density,

$$\mathcal{L} = -g_s Z'_\mu J^\mu, \quad (22)$$

where the current is given by

$$J^\mu = \bar{\psi} Q' \gamma^\mu \psi. \quad (23)$$

In the above equation, the summation of fermion fields is implied and the generator Q' is given approximately by

$$Q' \simeq \sqrt{\frac{3}{8}} \left(T + \frac{4}{9} \frac{g_L^2}{g_s^2} I_{3R} \right), \quad (24)$$

where we have again assumed that $\sin^2 \theta_w(M_{W'}) \simeq 1/4$. From the contributions of the $Z', W_R^\pm$ to low energy experiments, a limit of $M_{Z'}, M_{W_R} \gtrsim 0.5 - 1 \text{ TeV}$ is expected[10]. Thus, we argue that the model is quite weakly constrained given that the exotic symmetry breaking scale can be as low as a TeV or so.

The model contains scalar lepto-quarks which could be relatively light (e.g. a few hundred GeV). In particular χ_L contains $SU(3)_c$ triplet $SU(2)_L$ doublet scalars

coupling the left-handed leptons with the right handed d type quarks. From Eq.(4), we can deduce that

$$\mathcal{L}_\chi = \lambda_2 \bar{d}_R L_L \chi + H.c., \quad (25)$$

where χ is the colour triplet component of χ_L , and $L_L = (\nu_L, e_L)^T$. Note that χ -type lepto-quarks coupling to d_R with masses of around 200 GeV have been put forward as a possible explanation of the excess high Q^2 Hera events[11]. (However, the Hera anomaly is only a 2-3 sigma excess and may disappear when more data is taken).

Observe that the Yukawa Lagrangian density, Eq.(4) implies that the right-handed charged leptons will mix slightly with the E^- exotic fermions. For one generation, the mixing has the form,

$$\mathcal{L}_{mass} = (\bar{e}_L \bar{E}_L) \begin{pmatrix} \lambda_2 w_L & 0 \\ \lambda_3 u_1 + \lambda_4 u_2 & \lambda_1 w_R \end{pmatrix} \begin{pmatrix} e_R \\ E_R \end{pmatrix} + H.c. \quad (26)$$

Note however that this mixing is expected to be small because $u_{1,2} \ll w_R$ [given the bound Eq.(18)]. Because the exotic E^- fermions do not have canonical Z couplings, the mixing will induce small flavour changing neutral current (FCNC) couplings in the general case of three generation mixing. The mixing will be constrained by processes such as $\mu^- \rightarrow e^+ e^- e^-$.

Finally, I would like to comment briefly on a cosmological issue. Within the context of the standard big bang model, the phase transition at the temperature scale $T \sim w_R$ will generate monopoles. Monopoles occur because a semi-simple gauge theory is broken down to a gauge symmetry with a $U(1)$ factor. Monopoles in the early Universe can be a problem if they are too abundant. This issue has been examined in Ref.[7] for the case of the usual Pati-Salam model broken at a low scale. It was concluded that there is no problem if the symmetry breaking scale is low, which is the case that is being considered in the present paper.

In conclusion, an alternative Pati-Salam type gauge model has been proposed. The model allows quarks and leptons to be unified with gauge group $SU(4) \otimes SU(2)_L \otimes SU(2)_R$ at a relatively low scale. We argue that present data does not constrain this model very stringently (c.f. the usual Pati-Salam model). As a consequence the exotic gauge boson masses (and thus the symmetry breaking scale) can be as low as about 1 TeV. Neutrino masses arise radiatively in the model and are naturally light.

Acknowledgements

The author would like to thank Ray Volkas for the usual discussions and for pointing out some relevant papers. He would also like to thank X-G. He and J.P. Ma for a discussion and to J.Bowes for help with the Figure. The author is supported by an Australian Research Fellowship.

References

- [1] For a review, see e.g. V. Barger and R. J. N. Phillips, *Collider Physics*, (Addison-Wesley, 1987).
- [2] J. Pati and A. Salam, *Phys. Rev. D*10, 275 (1974).
- [3] R. Foot and H. Lew, *Phys. Rev. D*41, 3502 (1990).
- [4] R. Foot, H. Lew and R. R. Volkas, *Phys. Rev. D*44, 1531 (1991).
- [5] A. Kuznetsov and M. Mikheev, *Phys. Lett. B*329, 295 (1994).
- [6] G. Valencia and S. Willenbrock, *Phys. Rev. D*50, 6843 (1994).
- [7] R. R. Volkas, *Phys. Rev. D*53, 2681 (1996).
- [8] Particle data group, *Phys. Rev. D*54, 1 (1996).
- [9] See for example, J. L. Hewett et. al., *Argonne Accel. Phys.* 539 (1993). S. Davidson, D. Bailey and B. Campbell, *Z. Phys. C*61, 613 (1994); M. Leurer, *Phys. Rev. D*50, 536 (1994).
- [10] For a review, see for example, T. G. Rizzo, hep-ph/9501261 (1995).
- [11] See for example, J. L. Hewett and T. G. Rizzo, hep-ph/9703337 and references there-in.

Figure Caption

Figure 1: 1-loop Feynman diagram which leads to small electron neutrino Majorana mass. There will be similar diagrams for the other neutrinos. (The $W_L W_R$ mixing mass squared is obtained from $\mathcal{L} = (D_\mu \langle \phi \rangle)^\dagger D^\mu \langle \phi \rangle$ and is given by $\mu^2 = g_R g_L u_1 u_2$).

